

ACL + LATERAL KNEE RECONSTRUCTION PROTOCOL : APPROPRIATE FOR COMBINED ACL + LCL AND/OR PLC RECONSTRUCTIONS

PHASE 1

Monitor for evidence of:

- Infection: if patient develops a temperature $>38^{\circ}$, refer urgently to the operating surgeon. If the surgeon is unavailable, advise patient to attend A&E to exclude wound infection or septic arthritis
- Distal neurovascular deficit (*DVT, AECS, CPN* involvement)

Goals:

- Protect the grafts
- Control pain and swelling/effusion
- Restore/preserve range of motion
- **Muscle activation**
- Normal gait and movement patterns

Initial precautions:

- Avoid knee flexion $>90^{\circ}$ until **2 weeks**
- PWB (40% body weight) for **6 weeks** using long lever brace initially (locked at 0°)
- If able to SLR without extension lag, change long lever brace to Össur *CTi* brace at **2 weeks**
- Wean off crutches from **6 weeks** if normal gait, brace to be worn at all times for **12 weeks**
- Restricted OKC knee extension for **12 weeks** (more important for hamstrings ACL grafts)
- Avoid excessive hyperextension, external tibial rotation, tibial sag and varus postures for **16 weeks**
- Avoid open chain isolated hamstrings exercises for **16 weeks**

Pain, effusion and ROM:

- PEACE* protocol for the management of pain and swelling/effusion
NB: cryotherapy only influences pain, not drainage
- Passive/active assisted ROM $0-90^{\circ}$ for **2 weeks**, FROM thereafter (avoiding hyperextension)
- Patella mobilisation if required (medial/lateral, superior/inferior)

Muscle activation and strength:

- TAQ's, SLR in brace until able to perform without extension lag
- Double leg KKC ex's $\leq 70^{\circ}$ flexion (e.g. leg press) within weight bearing restrictions
- OKC knee extension with resistance between $90-45^{\circ}$ from **4 weeks**

Neuromuscular training:

- Double leg proprioceptive exercises (e.g. Bosu ball)

Cycling:

- Static bike with no resistance from **2 weeks** if sufficient ROM, increasing time as able

Criteria for progressing to Phase 2:

- Closed wound
- No/minimal pain with phase 1 exercises
- No/minimal synovitis/effusion
- Normal patellofemoral mobility, tibiofemoral ROM $\geq 0-120^{\circ}$
- Voluntary quadriceps contraction
- Normal FWB gait (from 6-8 weeks post-op)

AECS: Acute extremity compartment syndrome

CPN: Common peroneal nerve

CTI: Össur CTi (carbon titanium) brace

PEACE: Protection, Elevation, Avoid anti-inflammatories, Compression, Elevation.

ACL + LATERAL KNEE RECONSTRUCTION PROTOCOL – PHASE 2

Goals:

- Protect the grafts
- Full patellofemoral and tibiofemoral ROM
- **Correct movement patterns during exercises**
- Increase muscular endurance
- Protected lower limb strengthening
- **Wean off brace**

Precautions:

- Brace to be worn at all times until **12 weeks**, then wean off as able
- Restricted OKC knee extension until **12 weeks**
- Avoid excessive hyperextension, external tibial rotation, tibial sag and varus postures for **16 weeks**
- Avoid open chain isolated hamstrings exercises until **16 weeks**
- Avoid breaststroke, side stroke and whip kicking action in pool until **16 weeks**
- Avoid running until **20 weeks**

Strength:

- Double leg CKC ex's $\leq 70^\circ$, progress to single leg as able
- Double leg bridging from **week 10**
- Full range OKC knee extension with resistance from **12 weeks**
- Increase load on the quadriceps, gluteal and calf muscles.
- Start open chain isolated hamstrings exercises from **16 weeks**
- Progressively decrease repetitions and increase resistance for all strength exercises

Neuromuscular training:

- Increase difficulty of double leg proprioceptive ex's (e.g. perturbations, two motoric tasks)
- Control of knee varus and tibial external rotation at lower flexion angles ($<45^\circ$) during weight bearing exercises, using verbal, manual and visual cues as required
- Progress to single leg proprioceptive ex's as able

Cardiovascular exercises:

- Static bike with resistance from **9 weeks**
- Incline treadmill (7% gradient)
- Brisk walking programme over changing terrains as able
- Cross trainer or rower from **12 weeks**
- Stair/stepper machine from **16 weeks**
- Breaststroke, side stroke and whip kicking action in pool until **16 weeks**

Criteria for progressing to Phase 3:

- No/minimal pain with phase 2 exercises
- No/minimal synovitis/effusion
- Full ROM
- Correct qualitative performance of phase 2 exercise
- Successfully weaned off brace
- Able to walk briskly 3-5km over changing terrains without pain

ACL + LATERAL KNEE RECONSTRUCTION PROTOCOL – PHASE 3

Goals:

- Maintain good quality movement patterns
- Improve strength and power/rate of force development
- Increase difficulty of neuromuscular and perturbation training
- Start jogging and sports specific training

Precautions:

- Avoid excessive hyperextension, external tibial rotation, tibial sag and varus postures for **16 weeks**
- Avoid jogging/running until **20 weeks**
- Avoid functional testing (hop for distance, vertical hop, side hop) until **24 weeks**

Strength/power:

- Continue progressive loading for strengthening exercises
- Sports-specific progressions e.g. power development, jumping and landing.

Neuromuscular training:

- Increase difficulty of neuromuscular and perturbation training
- Emphasise sports specific movements
- Maintain quality of movement/performance during strength and sports exercises

Cardiovascular exercise:

- Increase intensity and duration of cardiovascular exercise
- Build sports specific load regarding energy expenditure (aerobic, anaerobic)

Running:

- Start running if:
 - full ROM
 - pain \leq 2 VAS and no effusion despite adequate loading
 - limb symmetry index (LSI) \geq 70% for quadriceps and hamstrings strength
- Graduated running programme: start with 4-minute walk, 1-minute run (4:1) for 20 minutes
Decrease walking time and increase running time by 1 minute each week (3:2, 2:3, 1:4, 0:5)
Patient should be able to run for 20 minutes after 5 weeks.
- Once running programme complete, introduce backwards and sideways running
- Progress running from single to multi-plane specific agility drills

Criteria for progressing to Phase 4:

- No/minimal pain with phase 3 rehabilitation
- Correct qualitative performance of phase 3 exercises
- Limb symmetry index (LSI) $>$ 80% for quads and hamstrings strength
- LSI $>$ 80% for hop battery tests (Gustavsson: hop for distance, vertical hop, side hop)

ACL + LATERAL KNEE RECONSTRUCTION PROTOCOL – PHASE 4

Goals:

- Sports specific drills and gradual return to play program
- Return to sport or physically demanding work

Precautions:

- Do not initiate progressive return to play programme until confirmation of restoration of lateral stability (<2mm side-side difference on varus stress X-ray) at **6 months** post-op

Strength/power:

- Sports-specific progressions e.g. power development, jumping and landing.

Neuromuscular training:

- Increase difficulty of neuromuscular and perturbation training (e.g. single legged jumps)
- Introduce reactive/unanticipated movements
- Emphasise sports specific movements
- Maintain quality of movement/performance during strength and sports exercises

Sports-specific training

- Increase intensity of agility training (e.g. cutting, pivoting)
- Build sports specific load regarding energy expenditure (aerobic, anaerobic)
- Build sports specific load regarding surface (grass, court etc.) and restart training with patient's team if confirmation of restoration of lateral stability at 6 months post-op

Criteria for returning to play:

- No knee pain with sports specific activities
- No giving way or fear of giving way during sports specific activities
- Active dynamic gait pattern and symmetrical jogging pattern
- Correct quality of performance with all sports-specific activities
- Limb symmetry index (LSI) >90% for quads and hamstrings strength
- LSI >90% for hop battery tests (Gustavsson: hop for distance, vertical hop, side hop)
- Drop test with analysis of movement (trunk, knee valgus and knee flexion when landing)
- Use ACL-RSI to measure patient's psychological readiness/confidence in return to sports
- Restoration of lateral stability confirmed by varus stress X-ray
- Minimal 9 months since surgery

Returning to sports >9 months post-op, and ensuring the patient has completed the return to sport criteria significantly reduces knee re-injury rate.

Isometric knee extensor torque values >3Nm/kg are associated with positive outcomes after ACLR.

Originator: Richard Norris, Orthopaedic Physiotherapy Specialist.

Ratified by: Mr M McNicholas, Consultant Orthopaedic Surgeon.

Date last reviewed: October, 2020

References:

ACL + LATERAL KNEE RECONSTRUCTION PROTOCOL – PHASE 1 APPROPRIATE FOR COMBINED ACL + LCL AND/OR PLC RECONSTRUCTIONS

Monitor for evidence of:

- Infection: if patient develops a temperature $>38^{\circ}$, refer urgently to the operating surgeon. If the surgeon is unavailable, advise patient to attend A&E to exclude wound infection or septic arthritis
- Distal neurovascular deficit (*DVT, AECS, CPN* involvement)

Goals:

- Protect the grafts
- Control pain and swelling/effusion
- Restore/preserve range of motion
- **Muscle activation**
- Normal gait and movement patterns

Initial precautions:

- Avoid knee flexion $>90^{\circ}$ until **2 weeks**
- PWB (40% body weight) for **6 weeks** using long lever brace initially (locked at 0°)
- If able to SLR without extension lag, change long lever brace to Össur *CTi* brace at **2 weeks**
- Wean off crutches from **6 weeks** if normal gait, brace to be worn at all times for **12 weeks**
- Restricted OKC knee extension for **12 weeks** (more important for hamstrings ACL grafts)
- Avoid excessive hyperextension, external tibial rotation, tibial sag and varus postures for **16 weeks**
- Avoid open chain isolated hamstrings exercises for **16 weeks**

Pain, effusion and ROM:

- PEACE* protocol for the management of pain and swelling/effusion
NB: cryotherapy only influences pain, not drainage
- Passive/active assisted ROM $0-90^{\circ}$ for **2 weeks**, FROM thereafter (avoiding hyperextension)
- Patella mobilisation if required (medial/lateral, superior/inferior)

Muscle activation and strength:

- TAQ's, SLR in brace until able to perform without extension lag
- Double leg CKC ex's $\leq 70^{\circ}$ flexion (e.g. leg press) within weight bearing restrictions
- OKC knee extension with resistance between $90-45^{\circ}$ from **4 weeks**

Neuromuscular training:

- Double leg proprioceptive exercises (e.g. Bosu ball)

Cycling:

- Static bike with no resistance from **2 weeks** if sufficient ROM, increasing time as able

Criteria for progressing to Phase 2:

- Closed wound
- No/minimal pain with phase 1 exercises
- No/minimal synovitis/effusion
- Normal patellofemoral mobility, tibiofemoral ROM $\geq 0-120^{\circ}$
- Voluntary quadriceps contraction
- Normal FWB gait (from 6-8 weeks post-op)

AECS: Acute extremity compartment syndrome

CPN: Common peroneal nerve

CTi: Össur CTi (carbon titanium) brace

PEACE: Protection, Elevation, Avoid anti-inflammatories, Compression, Elevation.

ACL + LATERAL KNEE RECONSTRUCTION PROTOCOL – PHASE 2

Goals:

- Protect the grafts
- Full patellofemoral and tibiofemoral ROM
- **Correct movement patterns during exercises**
- Increase muscular endurance
- Protected lower limb strengthening
- **Wean off brace**

Precautions:

- Brace to be worn at all times until **12 weeks**, then wean off as able
- Restricted OKC knee extension until **12 weeks**
- Avoid excessive hyperextension, external tibial rotation, tibial sag and varus postures for **16 weeks**
- Avoid open chain isolated hamstrings exercises until **16 weeks**
- Avoid breaststroke, side stroke and whip kicking action in pool until **16 weeks**
- Avoid running until **20 weeks**

Strength:

- Double leg CKC ex's $\leq 70^\circ$, progress to single leg as able
- Double leg bridging from **week 10**
- Full range OKC knee extension with resistance from **12 weeks**
- Increase load on the quadriceps, gluteal and calf muscles.
- Start open chain isolated hamstrings exercises from **16 weeks**
- Progressively decrease repetitions and increase resistance for all strength exercises

Neuromuscular training:

- Increase difficulty of double leg proprioceptive ex's (e.g. perturbations, two motoric tasks)
- Control of knee varus and tibial external rotation at lower flexion angles ($<45^\circ$) during weight bearing exercises, using verbal, manual and visual cues as required
- Progress to single leg proprioceptive ex's as able

Cardiovascular exercises:

- Static bike with resistance from **9 weeks**
- Incline treadmill (7% gradient)
- Brisk walking programme over changing terrains as able
- Cross trainer or rower from **12 weeks**
- Stair/stepper machine from **16 weeks**
- Breaststroke, side stroke and whip kicking action in pool until **16 weeks**

Criteria for progressing to Phase 3:

- No/minimal pain with phase 2 exercises
- No/minimal synovitis/effusion
- Full ROM
- Correct qualitative performance of phase 2 exercise
- Successfully weaned off brace
- Able to walk briskly 3-5km over changing terrains without pain

ACL + LATERAL KNEE RECONSTRUCTION PROTOCOL – PHASE 3

Goals:

- Maintain good quality movement patterns
- Improve strength and power/rate of force development
- Increase difficulty of neuromuscular and perturbation training
- Start jogging and sports specific training

Precautions:

- Avoid excessive hyperextension, external tibial rotation, tibial sag and varus postures for **16 weeks**
- Avoid jogging/running until **20 weeks**
- Avoid functional testing (hop for distance, vertical hop, side hop) until **24 weeks**

Strength/power:

- Continue progressive loading for strengthening exercises
- Sports-specific progressions e.g. power development, jumping and landing.

Neuromuscular training:

- Increase difficulty of neuromuscular and perturbation training
- Emphasise sports specific movements
- Maintain quality of movement/performance during strength and sports exercises

Cardiovascular exercise:

- Increase intensity and duration of cardiovascular exercise
- Build sports specific load regarding energy expenditure (aerobic, anaerobic)

Running:

- Start running if:
 - full ROM
 - pain ≤ 2 VAS and no effusion despite adequate loading
 - limb symmetry index (LSI) $\geq 70\%$ for quadriceps and hamstrings strength
- Graduated running programme: start with 4-minute walk, 1-minute run (4:1) for 20 minutes
Decrease walking time and increase running time by 1 minute each week (3:2, 2:3, 1:4, 0:5)
Patient should be able to run for 20 minutes after 5 weeks.
- Once running programme complete, introduce backwards and sideways running
- Progress running from single to multi-plane specific agility drills

Criteria for progressing to Phase 4:

- No/minimal pain with phase 3 rehabilitation
- Correct qualitative performance of phase 3 exercises
- Limb symmetry index (LSI) $>80\%$ for quads and hamstrings strength
- LSI $>80\%$ for hop battery tests (Gustavsson: hop for distance, vertical hop, side hop)

ACL + LATERAL KNEE RECONSTRUCTION PROTOCOL – PHASE 4

Goals:

- Sports specific drills and gradual return to play program
- Return to sport or physically demanding work

Precautions:

- Do not initiate progressive return to play programme until confirmation of restoration of lateral stability (<2mm side-side difference on varus stress X-ray) at **6 months** post-op

Strength/power:

- Sports-specific progressions e.g. power development, jumping and landing.

Neuromuscular training:

- Increase difficulty of neuromuscular and perturbation training (e.g. single legged jumps)
- Introduce reactive/unanticipated movements
- Emphasise sports specific movements
- Maintain quality of movement/performance during strength and sports exercises

Sports-specific training

- Increase intensity of agility training (e.g. cutting, pivoting)
- Build sports specific load regarding energy expenditure (aerobic, anaerobic)
- Build sports specific load regarding surface (grass, court etc.) and restart training with patient's team if confirmation of restoration of lateral stability at 6 months post-op

Criteria for returning to play:

- No knee pain with sports specific activities
- No giving way or fear of giving way during sports specific activities
- Active dynamic gait pattern and symmetrical jogging pattern
- Correct quality of performance with all sports-specific activities
- Limb symmetry index (LSI) >90% for quads and hamstrings strength
- LSI >90% for hop battery tests (Gustavsson: hop for distance, vertical hop, side hop)
- Drop test with analysis of movement (trunk, knee valgus and knee flexion when landing)
- Use ACL-RSI to measure patient's psychological readiness/confidence in return to sports
- Restoration of lateral stability confirmed by varus stress X-ray
- Minimal 9 months since surgery

Returning to sports >9 months post-op, and ensuring the patient has completed the return to sport criteria significantly reduces knee re-injury rate.

Isometric knee extensor torque values >3Nm/kg are associated with positive outcomes after ACLR.

Originator: Richard Norris, Orthopaedic Physiotherapy Specialist.

Ratified by: Mr M McNicholas, Consultant Orthopaedic Surgeon.

Date last reviewed: October, 2020

References:

1. UK National Ligament Registry: <https://www.uknlr.co.uk/im-a-patient.php>
2. Cartwright-Terry M, Yates J, Tan CK, Pengas IP, Banks JV, McNicholas MJ. Medium-Term (5 Year) Comparison of the Functional Outcomes of Combined Anterior Cruciate Ligament and Posterolateral Corner Reconstruction Compared with Isolated Anterior Cruciate Ligament Reconstruction. *Arthroscopy: The Journal of Arthroscopic and Related Surgery*, (2014). 30 (7), 811-817.
3. Development and Validation of a Short Version of the Anterior Cruciate Ligament Return to Sport After Injury (ACL-RSI) Scale. KE Webster JA Feller. *Orthop J Sports Med*. 2018 Apr; 6(4): 2325967118763763.
4. Dubois B, Esculier JF. Soft-tissue injuries simply need PEACE and LOVE. *Br J Sports Med*. 2020;54(2):72-3.
5. Imoto AM, Peccin S, Almeida GJ, Saconato H, Atallah Á. Effectiveness of electrical stimulation on rehabilitation after ligament and meniscal injuries: a systematic review. *Sao Paulo Med J*. 2011;129(6):414-23.
6. Grindem, Hege, et al. "Simple decision rules can reduce reinjury risk by 84% after ACL reconstruction: the Delaware-Oslo ACL cohort study." *Br J Sports Med* 50.13 (2016): 804-808.
7. Hauger AV, Reiman MP, Bjordal JM, Sheets C, Ledbetter L, Goode AP. Neuromuscular electrical stimulation is effective in strengthening the quadriceps muscle after anterior cruciate ligament surgery. *Knee Surg Sports Traumatol Arthrosc*. 2018;26(2):399-410.
8. Kuenze C, Hertel J, Saliba S, Diduch DR, Weltman A, Hart JM. Clinical thresholds for quadriceps assessment after anterior cruciate ligament reconstruction. *J Sport Rehabil*. 2015;24(1):36-46.
9. Kyritsis, Polyvios, et al. "Likelihood of ACL graft rupture: not meeting six clinical discharge criteria before return to sport is associated with a four times greater risk of rupture." *Br J Sports Med* (2016): bjsports-2015.
10. LaPrade, R.F., Wijdicks, C. A. (2012) The Management of Injuries to the Medial Side of the Knee, *Journal of Orthopaedic & Sports Physical Therapy*. Vol: 42 (3), pp 221-233.
11. LaPrade RF, Chahla J, DePhillipo NN, Cram T, Kennedy MI, Cinque M, et al. Single-Stage Multiple-Ligament Knee Reconstructions for Sports-Related Injuries: Outcomes in 194 Patients. *Am J Sports Med*. 2019;47(11):2563-71.
12. Perriman A, Leahy E, Semciw AI. The Effect of Open- Versus Closed-Kinetic-Chain Exercises on Anterior Tibial Laxity, Strength, and Function Following Anterior Cruciate Ligament Reconstruction: A Systematic Review and Meta-analysis. *J Orthop Sports Phys Ther*. 2018;48(7):552-66.
13. Pietrosimone B, Lepley AS, Harkey MS, Luc-Harkey BA, Blackburn JT, Gribble PA, et al. Quadriceps Strength Predicts Self-reported Function Post-ACL Reconstruction. *Med Sci Sports Exerc*. 2016;48(9):1671-7.
14. Rambaud AJM, Ardern CL, Thoreux P, Regnaud JP, Edouard P. Criteria for return to running after anterior cruciate ligament reconstruction: a scoping review. *Br J Sports Med*. 2018;52(22):1437-44.
15. van Grinsven, S., et al. "Evidence-based rehabilitation following anterior cruciate ligament reconstruction." *Knee Surgery, Sports Traumatology, Arthroscopy* 18.8 (2010): 1128-1144.
16. van Melick, Nicky, et al. "Evidence-based clinical practice update: practice guidelines for anterior cruciate ligament rehabilitation based on a systematic review and multidisciplinary consensus." *Br J Sports Med* (2016): bjsports-2015